Space Weather Highlights 22-28 December 1997

Solar activity was very low to low. Region 8124 (S22, L = 348, class/area EKC/590 on 23 December) grew at a moderate pace during 22 - 23 December and produced isolated B- and C-class subflares. The region's most notable flare was a brief C2/SF, reaching maximum at 26/0557UT, and accompanied by a partial-halo coronal mass ejection. There were no discrete or sweep frequency bursts detected with this flare.

Solar wind data were received from the WIND spacecraft a few hours per day. Velocities mostly ranged from 270 - 340 km/sec, but briefly increased to 400 km/sec on 23 December. Densities briefly increased to 31 p/cc on 23 December, but were in the 05 - 18 p/cc range for the rest of the period. Bz ranged from plus 05 to minus 05 nT (GSM). Solar sector orientation was mostly toward (phi angle near 315 degrees) during 26 - 27 December, but was not discernible during the rest of the period.

There were no significant proton enhancements observed at geosynchronous altitude.

The greater than 2 MeV electron flux at geosynchronous altitude was at normal levels.

The geomagnetic field was quiet at all latitudes.

Space Weather Forecast 31 December 1997 - 26 January 1998

Solar activity is expected to range from very low to low. Isolated C-class flare activity is expected. No significant proton enhancements are expected at geosynchronous altitude.

The greater than 2 MeV electron flux at geosynchronous altitude is expected to be at normal levels during most of the period.

The geomagnetic field is expected to be quiet to unsettled.



Daily Solar Data

| | Radio | Sun | Sunspot | X-ray | | | | Flares | | | | |
|-------------|---------|--------|-------------------------|------------|---------|---------|----|--------|---|---|---|---|
| | Flux | spot | Area | Background | X | -ray Fl | ux | | | | | |
| Date | 10.7 cm | No. (1 | 10 ⁻⁶ hemi.) | | C | M | X | S | 1 | 2 | 3 | 4 |
| 22 December | r 97 | 41 | 260 | B1.9 | 3 | 0 | 0 | 2 | 0 | 0 | 0 | 0 |
| 23 December | r 104 | 72 | 700 | B2.4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 24 December | r 108 | 67 | 520 | B2.2 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 |
| 25 December | r 105 | 56 | 470 | B1.6 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 |
| 26 December | r 105 | 50 | 460 | B2.1 | 1 | 0 | 0 | 2 | 0 | 0 | 0 | 0 |
| 27 December | r 96 | 56 | 420 | B1.6 | 1 | 0 | 0 | 0 | 1 | 0 | 0 | 0 |
| 28 December | r 102 | 55 | 410 | B1.4 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 |
| | | | | Daily Par | ticle T | ata | | | | | | |

Proton Fluence Electron Fluence (protons/cm²-day-sr) (electrons/cm²-day-sr) >.6MeV Date >1MeV >10MeV >100MeV >2MeV >4MeV 5.5E+5 22 December 2.6E+5 1.8E+44.2E+323 December 2.0E+5 1.7E+44.2E+32.9E+524 December 2.1E+5 1.7E+44.4E+34.8E+525 December 2.3E+5 1.8E+44.3E+36.9E + 526 December 1.8E+5 1.8E+44.5E+37.0E + 527 December 2.2E+5 1.8E+44.4E+37.7E+528 December 3.9E+5 1.8E+44.5E+38.3E+5

| | Daily Geomagnetic Data | | | | | | | | | | | | | |
|-------------|------------------------|-----------------|---|-----------------|---|-----------------|--|--|--|--|--|--|--|--|
| | Middle Latitude | | | High Latitude | I | Estimated | | | | | | | | |
| | F | Fredericksburg | | College | | Planetary | | | | | | | | |
| Date | A | K-indices | A | K-indices | A | K-indices | | | | | | | | |
| 22 December | 2 | 0-0-1-1-1-1-1 | 0 | 0-0-0-2-0-0-0 | 2 | 0-0-0-1-1-1-1 | | | | | | | | |
| 23 December | 3 | 1-1-0-0-1-1-1-2 | 0 | 0-0-0-0-0-0-0 | 3 | 1-0-0-0-1-1-1-2 | | | | | | | | |
| 24 December | 4 | 2-2-1-0-1-1-1 | 1 | 0-1-1-0-1-0-0-0 | 3 | 2-2-1-0-1-1-1-0 | | | | | | | | |
| 25 December | 2 | 1-1-1-0-0-0-1-0 | 0 | 0-0-0-0-0-0-0 | 2 | 1-1-0-0-0-0-1-0 | | | | | | | | |
| 26 December | 2 | 1-0-0-0-1-0-1-1 | 0 | 0-0-0-0-0-0-0 | 1 | 1-0-0-0-0-1-0 | | | | | | | | |
| 27 December | 0 | 0-0-0-0-0-1-0 | 0 | 0-0-0-0-0-0-0 | 2 | 0-0-0-0-0-1-0 | | | | | | | | |
| 28 December | 1 | 0-0-1-0-0-0-1-1 | * | 0-*-0-0-0-0-0 | 2 | 0-0-0-1-1-1-1 | | | | | | | | |




Alerts and Warnings Issued

| | <u> </u> | |
|---------------------------|--------------------------------------|-----------------------------|
| Date and Time of Issue (U | JT) Type of Alert or Warning | Date and Time of Event (UT) |
| 22 Dec 1355 | Stratwarm Alert exists Stratwarm M | onday |
| 23 Dec 1358 | Stratwarm Alert Exists Stratwarm Tu | ıesday |
| 24 Dec 1330 | Stratwarm Alert Exists Stratwarm Wed | dnesday |
| 25 Dec 1456 | Stratwarm Alert Exists Stratwarm Th | ursday |
| 26 Dec 1436 | Stratwarm Alert Exists Stratwarm F | riday |
| 27 Dec 1358 | Stratwarm Alert Exists Stratwarm Sa | turday |
| 28 Dec 1323 | Stratwarm Alert Exists Stratwarm St | unday |
| | | • |



Twenty-seven Day Outlook



| | Radio Flux | Planetary | Largest | | Radio Flux | Planetary | Largest |
|--------|------------|-----------|----------|--------|------------|-----------|----------|
| Date | 10.7 cm | A Index | Kp Index | Date | 10.7 cm | A Index | Kp Index |
| 31 Dec | 104 | 13 | 3 | 14 Jan | 86 | 5 | 2 |
| 01 Jan | 104 | 10 | 3 | 15 | 88 | 5 | 2 |
| 02 | 102 | 10 | 3 | 16 | 90 | 5 | 2 |
| 03 | 98 | 8 | 3 | 17 | 92 | 5 | 2 |
| 04 | 96 | 5 | 2 | 18 | 94 | 5 | 2 |
| 05 | 92 | 5 | 2 | 19 | 96 | 5 | 2 |
| 06 | 90 | 8 | 3 | 20 | 98 | 5 | 2 |
| 07 | 88 | 8 | 3 | 21 | 100 | 5 | 2 |
| 08 | 86 | 5 | 2 | 22 | 102 | 5 | 2 |
| 09 | 86 | 5 | 2 | 23 | 102 | 5 | 2 |
| 10 | 86 | 5 | 2 | 24 | 102 | 10 | 3 |
| 11 | 86 | 5 | 2 | 25 | 102 | 10 | 3 |
| 12 | 86 | 5 | 2 | 26 | 102 | 10 | 3 |
| 13 | 86 | 5 | 2 | | | | |



Energetic Events

| | Time (U | JT) | X-ray | Op | otical Informati | on | Peak | Sweep Freq |
|------|-----------|-----|------------|-------|------------------|-----|------------|------------|
| Date | | 1/2 | Integ | Imp | Location | Rgn | Radio Flux | Intensity |
| | Begin Max | Max | Class Flux | Brtns | Lat CMD | # | 245 2695 | II IV |

No Events Observed

| 777 | T . |
|-------|------|
| HIMPO | 1101 |
| | |

| | | | | Flare List | | | |
|-------------|-------------|-----------|------|------------|------------|----------|------|
| | | | | | | ptical | |
| ъ. | | Time | | X-ray | Imp / | Location | Rgn |
| Date | Begin | Max | End | Class. | Brtns | Lat CMD | # |
| | | | | | | | |
| 22 December | 0155 | 0220 | 0238 | C3.3 | | | |
| | 0622 | 0627 | 0636 | B3.8 | | | |
| | 0830 | 0830 | 0834 | C1.4 | SF | S24E64 | 8124 |
| | 1212 | 1217 | 1222 | B4.0 | | | |
| | 1238 | 1243 | 1247 | B4.3 | | | |
| | 1524 | 1536 | 1546 | C6.5 | SF | S24E58 | 8124 |
| | 2102 | 2108 | 2112 | B4.2 | | | |
| | 2151 | 2156 | 2200 | B7.1 | | | |
| | 2229 | 2235 | 2240 | B5.3 | | | |
| | 2245 | 2249 | 2254 | B5.6 | | | |
| 23 December | No Fla | res Obser | ved | | | | |
| 24 December | 0119 | 0122 | 0125 | B5.2 | | | |
| | 0302 | 0306 | 0315 | B3.0 | | | |
| | 1545 | 1550 | 1601 | B4.3 | | | |
| | 1611 | 1611 | 1616 | | SF | S22E25 | 8124 |
| | 1908 | 1908 | 1911 | B8.0 | SF | N24E72 | 8126 |
| 25 December | 0212 | 0213 | 0218 | B4.7 | SF | S22E23 | 8124 |
| | 0327 | 0330 | 0332 | B2.3 | | | |
| | 0955 | 0956 | 1000 | B3.2 | SF | S22E18 | 8124 |
| 26 December | 0409 | 0410 | 0418 | | SF | S24E14 | 8124 |
| | 0554 | 0554 | 0611 | C2.0 | SF | S22E07 | 8124 |
| 27 December | 1228 | 1238 | 1253 | B4.9 | | | |
| | 1632 | 1634 | 1704 | C3.2 | 1F | S23W06 | 8124 |
| 28 December | 1413 | 1417 | 1421 | B2.3 | | 2=2 30 | |
| 5 = 5555550 | 1751 | 1752 | 1756 | B5.0 | SF | S22W25 | 8124 |
| | 1,01 | 1,02 | 1,00 | 20.0 | ~ 1 | 522 25 | |
| | | | | | | | |



Region Summary

| | Location | | | Characteri | | | _ | | | Fl | ares | | | _ | |
|-------|---------------|---------|-------------------------|------------|-------|-------|-------|---|-------|----|------|---|-------|---|---|
| _ | (01 000) | Helio | Area | Extent | Spot | Spot | Mag | | X-ray | | _ | | ptica | | _ |
| Date | (° Lat ° CMD) | Lon | (10 ⁻⁶ hemi) | (helio) | Class | Count | Class | С | M | X | S | 1 | 2 | 3 | 4 |
| | Re | gion 81 | 23 | | | | | | | | | | | | |
| 14 De | ec N19E61 | 086 | 0010 | 04 | BXO | 003 | В | | | | | | | | |
| 15 De | c N18E46 | 088 | 0060 | 06 | DSO | 007 | В | 1 | | | 6 | | | | |
| 16 De | ec N18E32 | 088 | 0150 | 07 | DSO | 010 | В | | | | | | | | |
| 17 De | ec N18E20 | 087 | 0130 | 08 | DSO | 010 | В | | | | | | | | |
| 18 De | c N19E06 | 088 | 0100 | 09 | DSO | 014 | В | | | | | | | | |
| 19 De | ec N18W08 | 089 | 0070 | 10 | CSO | 006 | В | | | | 1 | | | | |
| | ec N18W20 | 088 | 0070 | 10 | DSO | 006 | В | | | | | | | | |
| | ec N18W36 | 091 | 0040 | 07 | CSO | 003 | В | | | | | | | | |
| 22 De | ec N18W51 | 092 | 0020 | 01 | HSX | 001 | A | | | | | | | | |
| 23 De | ec N18W64 | 092 | 0000 | 01 | AXX | 001 | A | | | | | | | | |
| 24 De | ec N18W77 | 092 | 0000 | 00 | AXX | 001 | A | | | | | | | | |
| 25 De | ec N18W90 | 092 | | | | | | | | | | | | | |
| | | | | | | | | 1 | 0 | 0 | 7 | 0 | 0 | 0 | 0 |
| Cross | ed West Lim | b. | | | | | | | | | | | | | |
| Absol | ute heliograp | hic lon | gitude: 08 | 8 | | | | | | | | | | | |
| | | | | | | | | | | | | | | | |
| | Re | gion 81 | '24 | | | | | | | | | | | | |
| 20 De | ec S22E78 | 350 | 0040 | 02 | HSX | 001 | Α | | | | | | | | |
| | ec S22E66 | 349 | 0150 | 12 | EAO | 007 | В | | | | 1 | | | | |
| 22 De | ec S22E54 | 347 | 0240 | 14 | EAC | 020 | BG | 2 | | | 2 | | | | |
| 23 De | ec S22E40 | 348 | 0590 | 13 | EKC | 027 | BGD | | | | | | | | |
| 24 De | ec S22E27 | 348 | 0360 | 13 | EKC | 023 | BGD | | | | 1 | | | | |
| 25 De | ec S22E14 | 348 | 0310 | 13 | EAI | 024 | BG | | | | 2 | | | | |
| 26 De | ec S22E01 | 348 | 0260 | 12 | ESI | 017 | BG | 1 | | | 2 | | | | |
| 27 De | ec S21W12 | 347 | 0230 | 11 | ESC | 020 | BG | 1 | | | | 1 | | | |
| 28 De | ec S20W24 | 345 | 0190 | 12 | EAI | 015 | BG | | | | 1 | | | | |
| | | | | | | | | 4 | 0 | 0 | 9 | 1 | 0 | 0 | 0 |

Still on Disk.

Absolute heliographic longitude: 348



Region Summary- continued.

| Location | | | | Sunspot | Characteri | stics | | | | | Fla | ares | | | |
|----------|----------------------|-----------------|-------------------------|---------|------------|-------|-------|---|-------|---|-----|------|-------|---|--------------|
| _ | | Helio | Area | Extent | Spot | Spot | Mag | | X-ray | | _ | | ptica | | _ |
| Date (| (° Lat ° CMD) | Lon | (10 ⁻⁶ hemi) | (helio) | Class | Count | Class | С | M | X | S | 1 | 2 | 3 | 4 |
| | Region | 8125 | | | | | | | | | | | | | |
| 23 Dec | N19E41 | 347 | 0010 | 03 | BXO | 003 | В | | | | | | | | |
| 24 Dec | N19E28 | 347 | 0000 | 03 | BXO | 002 | В | | | | | | | | |
| 25 Dec | N19E16 | 346 | 0000 | 00 | AXX | 001 | A | | | | | | | | |
| 26 Dec | N19E03 | 346 | | | | | | | | | | | | | |
| 27 Dec | N19W10 | 346 | | | | | | | | | | | | | |
| | | | | | | | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Died or | n Disk. | | | | | | | | | | | | | | |
| Absolu | ite heliograp | hic long | gitude: 34 | 6 | | | | | | | | | | | |
| | Da | gion 81. | 26 | | | | | | | | | | | | |
| 23 Dec | N20E79 | 309 31. | 0100 | 03 | ННХ | 001 | A | | | | | | | | |
| | N21E66 | 309 | 0160 | 03 | HHX | | A | | | | 1 | | | | |
| | N21E54 | 308 | 0160 | 03 | HKX | | A | | | | 1 | | | | |
| | N20E41 | 308 | 0200 | 03 | СНО | 002 | В | | | | | | | | |
| | N21E28 | 307 | 0180 | 03 | HSX | 003 | A | | | | | | | | |
| | N22E16 | 305 | 0210 | 05 | СНО | 008 | В | | | | | | | | |
| | | | | | | | | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 |
| Still on | Disk. | | | | | | | | | | | | | | |
| Absolu | ite heliograp | hic long | gitude: 30 | 5 | | | | | | | | | | | |
| | D | . 01 | 27 | | | | | | | | | | | | |
| 26 Dag | . Ke S17W54 | gion 81. 043 | 0000 | 00 | AXX | 001 | ٨ | | | | | | | | |
| | : S17W54 : S17W67 | 043 | 0000 | UU | АЛЛ | 001 | A | | | | | | | | |
| 27 Dec | 31/W0/ | 043 | | | | | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Died or | n Disk | | | | | | | U | U | U | U | U | U | U | U |
| | ite heliograp | hic lone | gitude: 04 | 3 | | | | | | | | | | | |
| 1 100010 | | | 5.00.00 | | | | | | | | | | | | |
| | Re | gion 81. | 28 | | | | | | | | | | | | |
| 27 Dec | N28W54 | 029 | 0010 | 03 | BXO | 003 | В | | | | | | | | |
| 28 Dec | N28W68 | 029 | 0010 | 02 | BXO | 002 | В | | | | | | | | |
| | | | | | | | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Still on | | | | | | | | | | | | | | | |
| Absolu | te heliograp | hic long | gitude: 02 | 9 | | | | | | | | | | | |




Recent Solar Indices (preliminary) of the observed monthly mean values

| | | | | | monthly i | mean value: | | | |
|----------------|----------|-------|------------|--------|-----------|--------------------|---------|------------------|-------|
| | | | ot Numbers | | | | io Flux | Geomagne | |
| | Observed | | Ratio | Smooth | | Penticton | Smooth | <u>Planetary</u> | |
| Month | SWO | RI | RI/SWO | SWO | RI | 10.7 cm | Value | Ap | Value |
| | | | | | 1995 | | | | |
| December | 16.2 | 10.0 | 0.62 | 17.6 | 10.8 | 72.6 | 72.8 | 09 | 10.0 |
| | | | | | 1996 | | | | |
| January | 17.6 | 11.5 | 0.65 | 16.8 | 10.4 | 74.5 | 72.4 | 09 | 09.8 |
| February | 09.1 | 04.4 | 0.48 | 16.2 | 10.1 | 71.5 | 72.2 | 10 | 09.8 |
| March | 12.1 | 09.2 | 0.76 | 15.4 | 09.7 | 72.7 | 72.1 | 11 | 09.9 |
| | | | | | | | | | |
| April | 08.5 | 04.8 | 0.56 | 13.6 | 08.5 | 69.3 | 71.6 | 11 | 09.7 |
| May | 11.8 | 05.5 | 0.47 | 12.9 | 08.0 | 72.1 | 71.4 | 07 | 09.5 |
| June | 18.8 | 11.8 | 0.63 | 13.5 | 08.5 | 69.6 | 71.8 | 05 | 09.4 |
| | | | | | | | | | |
| July | 13.2 | 08.2 | 0.62 | 13.4 | 08.4 | 71.2 | 72.0 | 07 | 09.3 |
| August | 20.5 | 14.4 | 0.70 | 13.1 | 08.3 | 72.4 | 72.1 | 09 | 09.4 |
| September | | 01.6 | 0.55 | 13.3 | 08.4 | 69.4 | 72.3 | 15 | 09.3 |
| z ep terrie er | 02.0 | 01.0 | 0.00 | 10.0 | 0011 | 0,,, | , | 10 | 0,10 |
| October | 02.3 | 00.9 | 0.39 | 14.0 | 08.8 | 69.2 | 72.6 | 13 | 09.1 |
| November | 26.7 | 17.9 | 0.67 | 15.4 | 09.8 | 78.7 | 73.0 | 08 | 09.1 |
| December | 21.1 | 13.3 | 0.63 | 16.2 | 10.4 | 77.8 | 73.3 | 07 | 09.3 |
| | | | | | 1997 | | | | |
| January | 09.0 | 05.7 | 0.63 | 16.5 | 10.5* | 74.0 | 73.4 | 09 | 09.3* |
| February | 11.3 | 07.6 | 0.67 | 17.4 | 11.0* | 73.8 | 73.7 | 11 | 09.2* |
| March | 14.4 | 08.7 | 0.60 | 20.4 | 13.5* | 73.5 | 75.1* | 08 | 09.0* |
| 1,141,411 | - | 00.7 | 0.00 | | 20.0 | , , , , | 7011 | | 0,10 |
| April | 24.5 | 15.5 | 0.63 | 24.0 | 16.5* | 74.5 | 76.8* | 10 | 08.6* |
| May | 28.6 | 18.5 | 0.65 | 26.4 | 18.4* | 74.6 | 78.4* | 08 | 08.6* |
| June | 22.1 | 12.7 | 0.57 | | | 71.7 | | 07 | |
| Julie | 22.1 | 12.7 | 0.57 | | | , 1., | | 07 | |
| July | 17.0 | 10.5* | 0.62* | | | 71.1 | | 06* | |
| August | 36.7 | 24.7* | 0.67* | | | 79.0 | | 08* | |
| September | | 51.3* | 0.88* | | | 96.2* | | 10* | |
| September | 22.0 | 31.3 | 3.00 | | |) U.L | | 10 | |
| October | 33.6 | 23.3* | 0.69* | | | 85.0* | | 10* | |
| November | 53.5 | 39.3* | 0.73* | | | 99.5* | | 10* | |
| | 22.2 | 27.0 | 00 | | | , , . . | | | |

^{*}Preliminary estimates.

The lowest smoothed sunspot number for Cycle 22, RI = 8.0, occurred May 1996. The highest smoothed sunspot number for Cycle 22, RI=158.5, occurred July 1989. October 1996 has been selected as the start of Solar Cycle 23. From June 1991 onward, the 10.7 cm radio flux data source is Penticton, B.C. Canada. Prior to that, it was Ottawa.





Weekly Geosynchronous Satellite Environment Summary

Week Beginning 22 December 1997


Protons plot contains the five-minute averaged integral proton flux (protons/ cm²-sec-sr) as measured by GOES-9 (W135) for each of three energy thresholds: greater than 10, 50, and 100 MeV.


Electrons plot contains the five-minute averaged integral electron flux (electrons/ cm² -sec-sr) with energies greater than 2 MeV at GOES-9.

Hp plot contains the five minute averaged magnetic field H component in nanoteslas (nT) as measured by GOES-9. The H component is parallel to the spin axis of the satellite, which is nearly parallel to the Earth's rotation axis. *Kp* plot contains the estimated planetary 3-hour K-index (derived by the USAF 55th Space Weather Squadron) in real time from magnetometers at Meanook, Canada; Sitka, AK; Glenlea, Canada; St. Johns, Canada; Ottawa, Canada; Newport, WA; Fredericksburg, VA; Boulder, CO; Fresno, CA and Heartland, UK. These data are made available through cooperation from the Geological Survey of Canada (GSC) and the US Geological Survey. These may differ from the final Kp values derived from a more extensive network of magnetometers.

The data included here are those now available in real time at the SWO and are incomplete in that they do not include the full set of parameters and energy ranges known to cause satellite operating anomalies. The proton and electron fluxes and Kp are "global" parameters that are applicable to a first order approximation over large areas. Hparallel is subject to a more localized phenomena and the measurements generally are applicable to within a few degrees of longitude of the measuring satellite.









Weekly GOES Satellite X-ray and Proton Plots


Proton plot contains the five minute averaged integral proton flux (protons/cm²-sec-sr) as measured by GOES-9 (W135) for each of the energy thresholds: >1, >10, >30 and >100 MeV. P10 event threshold is 10 pfu (protons/cm²-sec-sr) at greater than 10 MeV.


X-ray plot contains five minute averaged x-ray flux (watts/m²) as measured by GOES 8 and 9 in two wavelength bands, .05 -.4 and .1 - .8 nm. The letters A, B, C, M and X refer to x-ray event levels for the .1 - .8 nm band.











12/19/97 No Data


12/20/97 No Data

12/21/97 no data

Coronal hole maps from the National Solar Observatory, Kitt Peak, Arizona

These maps are reproductions of the coronal hole contours as derived from the Kitt Peak 1083 nm raw data images. These are preliminary data, where solid (and dashed) lines are used to indicate more (and less) confidence in an inference and are printed whenever they are available at SWO.





Coronal hole maps from the National Solar Observatory, Kitt Peak, Arizona
These maps are reproductions of the coronal hole contours as derived from the Kitt Peak 1083 nm raw data images. These are preliminary data, where solid (and dashed) lines are used to indicate more (and less) confidence in an inference and are printed whenever they are available at SWO.

